

CONCOMITANT USE OF OPIOID AND BENZODIAZEPINE - EDUCATIONAL INTERVENTION Mailing conducted February 2017 – July 2019

Background:

During the April 2016 DUR Board Meeting, MS-DUR reviewed the CDC Guidelines for Prescribing Opioids for Chronic Pain¹ and data regarding DOM's performance on each recommendation that could be addressed through DUR efforts. One recommendation addressed concomitant use of opioids and benzodiazepines.

CDC recommendation: Providers should avoid prescribing opioid pain medication for patients receiving benzodiazepines whenever possible.

The CDC report noted that benzodiazepines and opioids both cause central nervous system depression and can decrease respiratory drive. Concurrent use is likely to put patients at greater risk for potentially fatal overdose. The clinical evidence review did not address risks of benzodiazepine co-prescription among patients prescribed opioids. However, the contextual evidence review found evidence in epidemiologic series of concurrent benzodiazepine use in large proportions of opioid-related overdose deaths, and a case-cohort study found concurrent benzodiazepine prescription with opioid prescription to be associated with a near quadrupling of risk for overdose death compared with opioid prescription alone.²

Experts agreed that although there are circumstances when it might be appropriate to prescribe opioids to a patient receiving benzodiazepines (e.g., severe acute pain in a patient taking long-term, stable low-dose benzodiazepine therapy), clinicians should avoid prescribing opioids and benzodiazepines concurrently whenever possible.

Table 6 below was included in the April DUR Board Packet. The distribution of beneficiaries taking opioids by number of days concurrent with taking benzodiazepines is shown in Table 6. Overall, 5.3% of beneficiaries taking opioids were concurrently taking benzodiazepines. Although this is a small percentage, it represents 6,376 beneficiaries that might be at increased risk of overdose death.

¹ CDC Guideline for Prescribing Opioids for Chronic Pain — United States, 2016.

<http://www.cdc.gov/media/modules/dpk/2016/dpk-pod/rr6501e1er-ebook.pdf>.

² Park TW, Saitz R, Ganoczy D, Ilgen MA, Bohnert AS. Benzodiazepine prescribing patterns and deaths from drug overdose among US veterans receiving opioid analgesics: case-cohort study. *BMJ* 2015;350:h2698. <http://dx.doi.org/10.1136/bmj.h2698>

TABLE 6: Distribution of Beneficiaries Taking Opioids by Number of Days Concurrent Use of Opioid and Benzodiazepine (2015 - Beneficiaries with cancer diagnoses are excluded)								
Number of Days Concurrently Taking Opioid and Benzodiazepine	TOTAL (n = 120,158)		FFS (n = 26,014)		UnitedHealth Care (n = 46,135)		Magnolia (n = 48,009)	
0	113,782	94.7%	25,130	96.6%	43,497	94.3%	45,155	94.1%
1 - 10	2,287	1.9%	382	1.5%	896	1.9%	1,009	2.1%
11 - 31	1,710	1.4%	271	1.0%	677	1.5%	762	1.6%
32 - 62	1,024	0.9%	92	0.4%	447	1.0%	485	1.0%
63 +	1,355	1.1%	139	0.5%	618	1.3%	598	1.2%

* Distributions are significantly different among plans ($p < 0.001$).

NOTE: Encounter data for Magnolia are not complete for November - December 2015.

The following motions were made and passed by the DUR Board:

- a. Concomitant use of opioids and benzodiazepines should require a manual prior authorization (PA).
- b. MS-DUR should provide an educational mailing to providers prescribing concurrent use of benzodiazepines and opioids to inform them of the increased safety risks and highlight the CDC recommendation to avoid concomitant use.

The Division of Medicaid (DOM) has initiated changes in the electronic PA process to require a manual PA for any new concomitant use of opioids and benzodiazepines. MS-DUR has initiated this educational mailing to address ongoing concomitant use.

MAILING

Exception monitoring was run for to identify benzodiazepine prescriptions being filled that resulted in concomitant use with opioids for more than 60 days in last 90 days. Providers were prioritized for mailings each month based on the number of beneficiaries with exceptions. Provider letters included information about up to five (5) beneficiaries each month. A provider was not contacted about the same beneficiary more than once every three months but could receive a letter each month.

The attached letter template and response form were used in the mailings.

The following number of providers received letters each month.

Mon+E4:K35th/Year		Total Prescriptions Analyzed for Mailing	Number of Prescriptions Exceeding Criteria	Number of Beneficiaries Exceeding Criteria	Number of Prescribers Mailed	Number of Beneficiaries Addressed in Letters
Mailed	Analyzed					
Feb-17	Dec-16	36864	5882	2600	150	727
Mar-17	Jan-17	37277	6442	3059	150	539
Apr-17	Feb-17	37971	6841	3232	150	637
May-17	Mar-17	28291	4134	1965	150	334
Jun-17	Apr-17	37076	6937	3295	150	664
Jul-17	May-17	35407	5875	2790	150	447
Aug-17	Jun-17	38309	6873	3250	150	635
Sep-17	Jul-17	33738	5427	2604	150	409
Oct-17	Aug-17	33887	5056	2396	150	532
Nov-17	Sep-17	34483	5496	2601	150	451
Dec-17	Oct-17	27975	3961	1903	150	485
Jan-18	Nov-17	29864	4336	2070	150	380
Feb-18	Dec-17	28391	3893	1854	150	485
Mar-18	Jan-18	30200	4218	2009	150	368
Apr-18	Feb-18	26660	3469	1645	150	412
May-18	Mar-18	15051	1477	701	150	187
Jun-18	Apr-18	15934	1762	832	150	283
Jul-18	May-18	27676	3234	1532	150	323
Aug-18	Jun-18	29429	3624	1710	150	405
Sep-18	Jul-18	28805	3268	1551	150	292
Oct-18	Aug-18	27470	2766	1314	150	321
Nov-18	Sep-18	20572	2081	990	150	232
Dec-18	Oct-18	26570	2692	1272	150	338
Jan-19	Nov-18	26813	2641	1240	150	276
Feb-19	Dec-18	24077	2225	1055	150	267
Mar-19	Jan-19	26322	2396	1124	150	249
Apr-19	Feb-19	26382	2251	1060	150	252
May-19	Mar-19	23454	1934	921	150	229
Jun-19	Apr-19	25395	2078	965	388	645
Jul-19	May-19	23385	1691	801	234	373

IMPORTANT INFORMATION ABOUT CONCOMITANT PRESCRIBING OF OPIOIDS AND BENZODIAZEPINES

<date>

Dear Dr. <prov_name>,

Dear [PRESCRIBER'S NAME],

On August 1, 2019, the Division of Medicaid (DOM) will implement several new pharmacy claims system edits as recommended by the Drug Utilization Review (DUR) Board in response to the Centers for Disease Control and Prevention (CDC) Guidelines for Prescribing Opioids for Chronic Pain and per the Centers for Medicare and Medicaid Services (CMS) requirements¹. These changes will be applicable for beneficiaries in the fee for service (FFS) and Coordinated Access Network (CAN) plans.

WHY YOU ARE RECEIVING THIS LETTER

DOM is mailing all providers with patients identified as concomitantly receiving benzodiazepines and opioids. Your patient(s) listed below was identified as being prescribed a benzodiazepine and opioid concomitantly. On August 1, 2019, these patients will require a prior authorization (PA) to continue concomitant use. The PA form can be found at

<https://medicaid.ms.gov/providers/pharmacy/pharmacy-prior-authorization/>.

WHAT WE ASK OF YOU?

- Whenever possible do not prescribe benzodiazepines and opioids for use at the same time.
- When prescribing benzodiazepines or opioids, check the Prescription Monitoring Program to be sure the prescription will not result in concomitant use of the two products. You can access the Prescription Monitoring Program at: <https://mississippi.pmpaware.net/login>
- If you think it is clinically necessary for a patient to take a benzodiazepine and an opioid at the same time, you must complete and submit a PA request form and attest to the following:
 1. Concomitant opioid and benzodiazepine therapy is medically necessary.
 2. The prescriber has acknowledged that he/she has informed the beneficiary about the risks of

¹ The Centers for Medicare and Medicaid Services (CMS) requires that state Medicaid programs have drug utilization review safety edits for opioid refills and an automated claims review process to identify refills in excess of state limits, monitor concurrent prescribing of opioids and benzodiazepines, on or prior to October 1, 2019. This is one of the many Medicaid-related provisions specified in Section 1004 of the SUPPORT Act (H.R. 6, the Substance Use-Disorder Prevention that Promotes Opioid Recovery and Treatment (SUPPORT) for Patients and Communities Act, the bipartisan bill aimed at addressing the nation's opioid overdose epidemic).

Evidence-Based DUR Initiative

concomitant utilization of opioid and benzodiazepine therapy or other drugs that could potentially cause respiratory depression and the beneficiary expressed understanding of these risks.

To obtain more information about the CDC Guidelines for Prescribing Opioids for Chronic Pain or to obtain provider resources related to the use of opioids, please use the link below:

<http://www.cdc.gov/drugoverdose/prescribing/resources.html>


Attached is a response form that you can use to provide us feedback about these beneficiaries. Should you have any questions or other comments about this intervention letter, please feel free to call MS-DUR at 662-915-7650.

Beneficiary Name	DOB	Opioid			Benzodiazepine		
		Drug Name	Date Filled	Prescriber	Drug Name	Date Filled	Prescriber
«Bene_name_1»	«bene_dob_1»	«drug_name_opi_1»	«fill_dt_opi_1»	«provi_name_opi_1»	«drug_name_benzo_1»	«fill_dt_benzo_1»	«provi_name_benzo_1»
«Bene_name_2»	«bene_dob_2»	«drug_name_opi_2»	«fill_dt_opi_2»	«provi_name_opi_2»	«drug_name_benzo_2»	«fill_dt_benzo_2»	«provi_name_benzo_2»
«Bene_name_3»	«bene_dob_3»	«drug_name_opi_3»	«fill_dt_opi_3»	«provi_name_opi_3»	«drug_name_benzo_3»	«fill_dt_benzo_3»	«provi_name_benzo_3»
«Bene_name_4»	«bene_dob_4»	«drug_name_opi_4»	«fill_dt_opi_4»	«provi_name_opi_4»	«drug_name_benzo_4»	«fill_dt_benzo_4»	«provi_name_benzo_4»
«Bene_name_5»	«bene_dob_5»	«drug_name_opi_5»	«fill_dt_opi_5»	«provi_name_opi_5»	«drug_name_benzo_5»	«fill_dt_benzo_5»	«provi_name_benzo_5»


Sincerely,



Carlos A. Latorre, MD, FAAFP
Medical Director
Mississippi Division of Medicaid



Terri R. Kirby, RPh, CPM
Director, Office of Pharmacy
Mississippi Division of Medicaid



Eric Pittman, PharmD
Project Director
MS-DUR



School of Pharmacy · University, MS 38677
phone: 662-915-7650 · fax: 662-915-5262
<http://www.pharmacy.olemiss.edu/cpm/m/medur.html>



Evidence-Based DUR Initiative

The Mississippi Division of Medicaid Drug Utilization Review Contractor



IMPORTANT INFORMATION ABOUT CONCOMITANT PRESCRIBING OF OPIOIDS AND BENZODIAZEPINES AND INCREASED RISK OF OVERDOSE AND DEATH

MD_NAME

Date of Mailing: DATE

Beneficiary Name	DOB	Opioid			Benzodiazepine		
		Drug Name	Date Filled	Prescriber	Drug Name	Date Filled	Prescriber
BENEFICIARY 1							
BENEFICIARY 2							
Up to 5 benes							

COMMENTS AND FEEDBACK:

Please check all that apply and add any additional comments or explanations desired

	BENEFICIARY 1	BENEFICIARY 2	Up to 5 benes		
My records do not indicate I wrote this(these) prescription(s)	<input type="checkbox"/>				
I was -					
• not aware of both prescriptions	<input type="checkbox"/>				
• aware of both prescriptions	<input type="checkbox"/>				
I consider the benefits of concomitant use to be greater than the risk for this patient.	<input type="checkbox"/>				
The patient has been counseled on the risks and benefits of concomitant use of these products.	<input type="checkbox"/>				
Additional comments:					

Please fax/mail your response to the number/address below.



School of Pharmacy • University, MS 38677
phone:662-915-7650 • fax:662-915-5262
<http://www.pharmacy.olemiss.edu/cpmm/msdur.html>